

# Carcinoid Tumors of the Colon and Rectum

P. KENT CULLEN, JR., MD  
Santa Barbara

IN 1808 MERLING first cited the carcinoid as an appendiceal tumor.<sup>1</sup> Five to ten percent of all carcinoid tumors are located in the colon, with the majority occurring in the cecum.<sup>2,3</sup> This tumor is the most common neoplasm of the appendix and is most often found in this organ.<sup>2,4,5</sup> The tumor is present in approximately 0.3 percent<sup>2</sup> of all appendices, variations depending upon the method of reporting. The literature is not clear on carcinoid tumors of "the colon" as reports are related to the appendix, rectum and colon or the colon exclusive of the rectum.<sup>6-9</sup>

## Age and Sex

As with other types of tumors, the incidence of carcinoid is highest in middle and old age. Appendicular carcinoid tumors usually occur at 25 to 30 years.<sup>10</sup> The average age of all patients with rectal carcinoids is 55 years.<sup>7</sup> Thirty-nine years is the mean age of patients with malignant rectal carcinoids.<sup>10</sup>

Considering all carcinoid tumors, there is no apparent difference in the incidence in males and females.<sup>10</sup> However, the incidence of carcinoids of the appendix alone is approximately twice as great in females as in males.<sup>10</sup>

## Symptoms

Carcinoids of the appendix are usually incidental findings, but occasionally they cause obstruction and are the direct cause of appendicitis. Carcinoids of the colon tend to be large, and the

symptoms are similar to those of carcinoma except that carcinoids have less tendency to ulcerate and bleed.<sup>4</sup> Colonic carcinoids tend to be large, the size correlating well with the high rate of metastasis.<sup>9</sup> Most rectal carcinoids produce few symptoms.<sup>6-8,11</sup> Rectal bleeding, change in bowel habits, varying degrees of obstruction, or signs and symptoms of metastatic carcinoma may be present.

## Pathology

A yellow character is often noted on cut section. The tumors are usually submucosal. Ulceration is not common, and necrosis rare.<sup>12</sup>

In tumors of the mid-gut, cytoplasmic granules can be demonstrated by the argentaffin reaction.<sup>13</sup> In lesions of the fore-gut and hind-gut, however, silver staining is variable.<sup>13</sup>

The incidence of metastasis from appendiceal lesions is only 3 percent because so often the appendix is removed routinely before dissemination can occur,<sup>4</sup> but from carcinoid tumors of the cecum and colon it is high.<sup>2,4,9</sup> Size is the best criterion by which to predict the malignant potential of these tumors in the rectum.<sup>6,7</sup> Lesions less than 1 cm in diameter are almost invariably benign, and those larger than 2 cm almost always develop local extension or metastasis. In addition to size, invasion into or beyond the true muscularis is an indication of malignancy. For that reason it is imperative that biopsy specimens be deep enough so that invasion can be seen.<sup>6,7</sup>

The reported incidence of multiple carcinoids of the colon<sup>9</sup> and rectum<sup>7</sup> has varied from 2 to 4.5 percent. For other malignant neoplasms associated with carcinoid tumors of the colon<sup>9</sup> and rectum<sup>7</sup> the range is from 2.5 to 38 percent.<sup>25</sup>

## Carcinoid Syndrome

Association of the carcinoid syndrome with appendiceal or colonic carcinoids is unusual.<sup>2</sup> There has been only one report of this syndrome in conjunction with a rectal lesion.<sup>14</sup> Ahlgren et al in 1962 published a case of carcinoid tumor of the rectum with moderate rise in serotonin without association of the carcinoid syndrome, the only case of a secreting carcinoid of the rectum described in the literature.<sup>15</sup> Orloff<sup>7</sup> accepts neither of those reports and presents his case well. Ingestion of serotonin-containing foods, such as bananas, tomatoes, avocados, red plums, walnuts and eggplant,<sup>16</sup> and drugs such as methocarbamol and mephensin carbamate,<sup>17</sup> elevate the urinary 5-hydroxyindoleacetic acid (5-HIAA) levels.

Presented at the Los Angeles County Division American Cancer Society Symposium, Neoplasms of the Colon and Rectum, St. Vincent's Hospital, February 14, 1973.

Reprint requests to: P. K. Cullen, Jr., MD, 230 W. Pueblo, Santa Barbara, CA 93105.

## Treatment

The treatment of carcinoid tumors is primarily surgical. When lymphatic permeation<sup>18-20</sup> or tumor at the surgical margin<sup>21</sup> is present, right hemicolectomy is advocated for appendiceal carcinoids. Considering that the prognosis of carcinoids of the appendix is excellent, with only a few deaths reported as directly attributable to the tumor,<sup>22</sup> one must be suspect of retreatment of this lesion regardless of the pathologic findings if the tumor is less than 2 cm in diameter.

In a review of the literature, Berardi<sup>9</sup> noted that the procedure most frequently used for colonic carcinoids was hemicolectomy. Anterior resection is the treatment of choice for sigmoid carcinoids.<sup>9</sup> When a nonresectable tumor was found, palliative colostomy or ileocolostomy was done.<sup>9</sup> In patients with the carcinoid syndrome (widespread metastatic disease) palliative resections to decrease the mass of tumor significantly improved the condition of some patients and prolonged life.<sup>9</sup> As was previously noted, recurrent carcinoids are extremely rare.<sup>23</sup>

Rectal carcinoids less than 2 cm in diameter, without muscle invasion, can be treated by local excision, excisional removal, and fulguration of the bed.<sup>6-8</sup> Larger tumors and those invading the muscle require radical extirpation, usually consisting of an abdominoperineal resection with permanent colostomy.<sup>6-8</sup>

Radiotherapy has proved to be of little value in the treatment of carcinoids except in an occasional case of metastasis to the liver.<sup>24</sup>

## Prognosis

The prognosis of carcinoid of the appendix is excellent. Moertel and coworkers<sup>5</sup> reported that in a series of 108 patients traced for more than five years and 83 for more than ten years, there were no recurrences regardless of the degree of local invasion. The same observers reviewed

more than a thousand cases in the literature and were able to find only two instances of recurrence.

Approximately one-fourth of all patients with malignant rectal carcinoids die postoperatively or from disease within five years.<sup>7</sup>

## REFERENCES

1. Merling: Cited in Forbus WD: Argentaffin tumors of the appendix and small intestine. *Bull Johns Hopkins Hosp.* 37:130-153, Aug 1925
2. Kultschitsky: Cited in Cheek RC, Wilson H: Carcinoid tumors. *Curr Prob Surg.* Chicago, Year Book Med Pub, Nov 1970, pp 4-31
3. Kuiper DH, Gracie WA, Pollard HH: Twenty years of gastrointestinal carcinoids. *Cancer* 25:1424-1430, Jun 1970
4. Sanders RJ, Axtell HK: Carcinoids of the gastrointestinal tract—A collective review. *Surg Gynec Obstet* 119:369-380, Aug 1964
5. Moertel CG, Dockerty MB, Judd ES: Carcinoid tumors of the vermiform appendix. *Cancer* 21:270-278, Feb 1968
6. Ponka JL, Walke L: Carcinoid tumors of the rectum. *Dis Col and Rect* 14:46-56, Jan-Feb 1971
7. Orloff, MJ: Carcinoid tumors of the rectum. *Cancer* 28:175-180, Jul 1971
8. Tiedemann RN, McDivitt RM, Thorbjarnarson B: Carcinoid tumors of rectum. *N Y State J Med* 72:559-563, Mar 1 1972
9. Berardi RS: Carcinoid tumors of the colon (exclusive of the rectum)—Review of the literature. *Dis Col and Rectum* 15:383-391, Sep-Oct 1972
10. Davies AJ: Carcinoid tumors (argentaffinomata). *Ann Roy Coll Surg Eng* 25:277-297, Dec 1959
11. Calderola VT, Jackman RJ, Moertel CG et al: Carcinoid tumors of the rectum. *Am J Surg* 107:844-849, Jun 1964
12. Teitelbaum SL: The carcinoid—A collective review. *Am J Surg* 123:564-572, May 1972
13. Williams ED, Sandler M: The classification of carcinoid tumors. *Lancet* 1:238-239, Feb 2, 1963
14. Saegesser F, Gross M: Carcinoid syndrome and carcinoid tumors of the rectum. *Am J. Proctol* 20:27-32, Feb 1969
15. Ahlgren M, Wirklander O: Urinary 5-HIAA and treatment of carcinoid of rectum—Report of one case. *Acta Chir Scand* 124:461-466, Nov 1962
16. Page IH: Serotonin. Chicago, Year Book Med Pub, 1968
17. Honet JC, Casey TV, Runyan JW: False-positive urinary test for 5-HIAA due to methocarbamol and mephensin carbamate. *N Engl J Med* 261:188-190, Jul 23, 1959
18. Farringer JL, Tarasidis G: Carcinoid tumors of the appendix. *Arch Surg* 88:354-356, Mar 1964
19. Campanale RP, Prothero SR: Malignant appendiceal carcinoid (argentaffin) with regional lymph node metastasis. *Am J Surg* 107:855-857, Jun 1964
20. Kieraldo J, Eversole S, Allen R: Carcinoid tumor of the vermiform appendix with distant metastasis—A review of the literature and report of two cases, one in a 14 year old girl. *Calif Med* 99:161-165, Sep 1963
21. Kantor S, Crane RD, Gillesby WJ: Carcinoid tumors of the gastrointestinal tract. *Am Surg* 27:448-453, Jun 1961
22. Turner M, Gallagher HS: Occult appendiceal carcinoid—Report of a case with fatal metastases. *Arch Path* 88:188-190, Aug 1969
23. Armstrong RB, Nielsen OF: Recurrent carcinoid tumor of the colon. *Arch Intern Med* 114:684-687, Nov 1964
24. Bagen JA: The carcinoid syndrome—Its diagnosis and management. *South Med J* 54:902-905, Aug 1961
25. Genre CF, Roth LM, Reed RJ: "Benign" rectal carcinoids—A report of two patients with metastases to regional lymph nodes. *Am J Clin Pathol* 56:750-757, Dec 1971